Remote Sensing of Lightning Impacts at the Edge of Space

Caitano L. da Silva1

¹Department of Physics & Langmuir Lab, New Mexico Tech, Socorro, NM, USA 87801

Abstract: The lower ionosphere marks the separation between neutral and ionized regions of the Earth's atmosphere, and has been colloquially referred to as the *edge of space*. The Earth and the ionosphere create a cavity that traps low-frequency electromagnetic waves, making long-range radio communications possible. Leveraging this concept, the high variability of the lower ionosphere, the so-called ionospheric D region, has been extensively probed with ELF/VLF remote sensing. A byproduct of thunderstorms, transient luminous events (TLEs), have appeared as an opportunity for optical remote-sensing of the lower ionosphere regions, but also its thermometer. In this presentation, we review our team's recent efforts to probe the impacts of lightning in the lower ionosphere by remote sensing of sprites and their electromagnetic signatures [Contreras-Vidal et al., 2021], and by looking at direct electron density changes measured by the Arecibo Observatory Incoherent Scatter Radar in coincidence with underlying thunderstorm activity [da Silva et al., 2021].

Acknowledgment: We thank the support from NSF via award AGS-1917069.

References:

Contreras-Vidal, L., R. G. Sonnenfeld, C. L. da Silva, M. McHarg, D. Jensen, J. Harley, L. Taylor, R. Haaland, and H. Stenbaek-Nielsen (2021), Relationship between sprite current and morphology, *J. Geophys. Res. Space Phys.*, 126 (3), e2020JA028930, doi: 10.1029/2020JA028930.

da Silva, C. L., S. D. Salazar, C. G. M. Brum, and P. Terra (2021), Survey of electron density changes in the daytime ionosphere over the Arecibo Observatory due to lightning and solar flares, *Scientific Reports*, 11(10250), doi: 10.1038/s41598-021-89662-x.

Session: Ionosphere and high atmosphere Oral or Poster: Oral